



TITLE:

Approximation of Expectation of Diffusion Processes based on Lie Algebra and Malliavin Calculus (Mathematical Economics)

AUTHOR(S):

Kusuoka, Shigeo

CITATION:

Kusuoka, Shigeo. Approximation of Expectation of Diffusion Processes based on Lie Algebra and Malliavin Calculus (Mathematical Economics). 数理解析研究所講究録 2003, 1337: 205-209

ISSUE DATE:

2003-08

URL:

<http://hdl.handle.net/2433/43394>

RIGHT:

Approximation of Expectation of Diffusion Processes based on Lie Algebra and Malliavin Calculus

Shigeo KUSUOKA

Graduate School of Mathematical Sciences

The University of Tokyo

In the present paper, we refine the idea in [1] by using notions in [5]. We use the notation in [5] for free Lie algebra. Let (Ω, \mathcal{F}, P) be a probability space and let $\{(B^1(t), \dots, B^d(t); t \in [0, \infty))\}$ be a d -dimensional Brownian motion. Let $B^0(t) = t$, $t \in [0, \infty)$. Let $V_0, V_1, \dots, V_d \in C_b^\infty(\mathbf{R}^N; \mathbf{R}^N)$. Here $C_b^\infty(\mathbf{R}^N; \mathbf{R}^n)$ denotes the space of \mathbf{R}^n -valued smooth functions defined in \mathbf{R}^N whose derivatives of any order are bounded. We regard elements in $C_b^\infty(\mathbf{R}^N; \mathbf{R}^N)$ as vector fields on \mathbf{R}^N .

Now let $X(t, x)$, $t \in [0, \infty)$, $x \in \mathbf{R}^N$, be the solution to the Stratonovich stochastic integral equation

$$X(t, x) = x + \sum_{i=0}^d \int_0^t V_i(X(s, x)) \circ dB^i(s). \quad (1)$$

Then there is a unique solution to this equation. Moreover we may assume that with probability one $X(t, x)$ is continuous in t and smooth in x .

Let $A = A_d = \{v_0, v_1, \dots, v_d\}$, be an alphabet, a set of letters, and A^* be the set of words consisting of A including the empty word which is denoted by 1. For $u = u^1 \dots u^k \in A^*$, $u^j \in A$, $j = 1, \dots, k$, $k \geq 0$, we denote by $n_i(u)$, $i = 0, \dots, d$, the cardinal of $\{j \in \{1, \dots, k\}; u^j = v_i\}$. Let $|u| = n_0(u) + \dots + n_d(u)$, a length of u , $\|u\| = |u| + n_0(u)$, and $\#(u)$ denotes the cardinal of $\{i \in \{0, \dots, d\}; n_i(u) \geq 1\}$ for $u \in A^*$. Let $\mathbf{R}\langle A \rangle$ be the \mathbf{R} -algebra of noncommutative polynomials on A , $\mathbf{R}\langle\langle A \rangle\rangle$ be the \mathbf{R} -algebra of noncommutative formal series on A , $\mathcal{L}(A)$ be the free Lie algebra over \mathbf{R} on the set A , and $\mathcal{L}((A))$ be the \mathbf{R} Lie algebra of free Lie series on the set A .

Let ι denotes the left normed bracketing operator, i.e.,

$$\iota(v_{i_1} \dots v_{i_n}) = [\dots [v_{i_1}, v_{i_2}], \dots, v_{i_n}].$$

Let $p : \mathbf{R}\langle A \rangle \rightarrow \mathbf{R}[x_0, \dots, x_d]$ denotes a natural homomorphism from the algebra of noncommutative polynomials to the the algebra of commutative polynomials such that $p(u) = x_0^{n_0(u)} \dots x_d^{n_d(u)}$, $u \in A^*$.

Vector fields V_0, V_1, \dots, V_d can be regarded as first differential operators over \mathbf{R}^N . Let $\mathcal{DO}(\mathbf{R}^N)$ denotes the set of smooth differential operators over \mathbf{R}^N . Then $\mathcal{DO}(\mathbf{R}^N)$ is a noncommutative algebra over \mathbf{R} . Let $\Phi : \mathbf{R}\langle A \rangle \rightarrow \mathcal{DF}(\mathbf{R}^N)$ be a homomorphism given by

$$\Phi(1) = \text{Identity}, \quad \Phi(v_{i_1} \dots v_{i_n}) = V_{i_1} \dots V_{i_n}, \quad n \geq 1, \quad i_1, \dots, i_n = 0, 1, \dots, d.$$

Also, note that

$$\Phi(\iota(v_{i_1} \cdots v_{i_n})) = [\cdots [V_{i_1}, V_{i_2}], \cdots, V_{i_n}], \quad n \geq 2, \quad i_1, \dots, i_n = 0, 1, \dots, d.$$

Let $B(t; u)$, $t \in [0, \infty)$, $u \in A^*$, be inductively defined by

$$B(t; 1) = 1, B(t; v_i) = B^i(t), \quad i = 0, 1, \dots, d,$$

and

$$B(t; uv_i) = \int_0^t B(s; u) \circ dB^i(s) \quad u \in A^*, \quad i = 0, \dots, d.$$

Also we define $B(t; w)$ $t \in [0, \infty)$, $w \in \mathbf{R}\langle A \rangle$ by

$$B(t; \sum_{u \in A^*} a_u u) = \sum_{u \in A^*} a_u B(t; u),$$

and we denote $B(1; w)$ by $B(w)$ for $w \in \mathbf{R}\langle A \rangle$.

Let $A_m^* = \{u \in A^*; \|u\| = m\}$, $m \geq 0$, and let $\mathbf{R}\langle A \rangle_m = \sum_{u \in A_m^*} \mathbf{R}u$, and $\mathbf{R}\langle A \rangle_{\leq m} = \sum_{k=0}^m \mathbf{R}\langle A \rangle_k$, $m \geq 0$. Let $j_m : \mathbf{R}\langle A \rangle \rightarrow \mathbf{R}\langle A \rangle_{\leq m}$ be a natural surjective linear map such that $j_m(u) = u$, $u \in A^*$, $\|u\| \leq m$, and $j_m(u) = 0$, $u \in A^*$, $\|u\| \geq m+1$. Let $\mathcal{L}(A)_m = \mathcal{L}(A) \cap \mathbf{R}\langle A \rangle_m$, and $\mathcal{L}(A)_{\leq m} = \mathcal{L}(A) \cap \mathbf{R}\langle A \rangle_{\leq m}$, $m \geq 1$. Let $A^{**} = \{u \in A^*; u \neq 1, v_0\}$, and $A_{\leq m}^{**} = \{u \in A^{**}; \|u\| \leq m\}$, $m \geq 1$.

Let $\Psi_s : \mathbf{R}\langle A \rangle \rightarrow \mathbf{R}\langle A \rangle$, $s > 0$, be given by

$$\Psi_s(\sum_{m=0}^{\infty} x_m) = \sum_{m=0}^{\infty} s^{m/2} x_m, \quad x_m \in \mathbf{R}\langle A \rangle_m, \quad m \geq 0.$$

Now we introduce a condition (UFG) on the family of vector field $\{V_0, V_1, \dots, V_d\}$ as follows.

(UFG) There are an integer ℓ and $\varphi_{u,u'} \in C_b^\infty(\mathbf{R}^N)$, $u \in A^{**}$, $u' \in A_{\leq \ell}^{**}$, satisfying the following.

$$\Phi(\iota(u)) = \sum_{u' \in A_{\leq \ell}^{**}} \varphi_{u,u'} \Phi(\iota(u')), \quad u \in A^{**}.$$

Let us define a semi-norm $\|\cdot\|_{V,n}$, $n \geq 1$, on $C_0^\infty(\mathbf{R}^N; \mathbf{R})$ by

$$\|f\|_{V,n} = \sum_{k=1}^n \sum_{u_1, \dots, u_k \in A^{**}, \|u_1 \cdots u_k\| = n} \|\Phi(\iota(u_1) \cdots \iota(u_k))f\|_\infty.$$

Now let us define a semigroup of linear operators $\{P_t\}_{t \in [0, \infty)}$ by

$$(P_t f)(x) = E[f(X(t, x))], \quad t \in [0, \infty), \quad f \in C_b^\infty(\mathbf{R}^N).$$

Then we can prove the following ([2]).

Theorem 1 Assume that the family of vector fields satisfies the condition (UFG). Then for any $n \geq 1$ there is a constant $C > 0$ such that

$$\|P_t f\|_{V,n} \leq \frac{C}{t^{n/2}} \|f\|_\infty, \quad f \in C_b^\infty(\mathbf{R}^N), \quad t \in (0, 1].$$

Let us think of a family $\{Q_{(s)}; s \in (0, 1]\}$ of linear operators in $C_b(\mathbf{R}^N)$.

Definition 2 We say that $Q_{(s)}$, $s \in (0, 1]$, is m -similar, $m \geq 1$, if there are a constant $C > 0$ and $n \geq m + 1$ such that

$$\|P_s f - Q_{(s)} f(x)\|_\infty \leq C \left(\sum_{k=m+1}^n s^{k/2} \|f\|_{V,k} + s^{(m+1)/2} \|\nabla f\|_\infty \right),$$

and

$$\|Q_{(s)} f - f\|_\infty \leq C s^{1/2} \|\nabla f\|_\infty$$

for any $s \in (0, 1]$, and $f \in C_b^\infty(\mathbf{R}^N; \mathbf{R})$.

Let $T > 0$ and $\gamma > 0$. Let $t_k = t_k^{(n)} = \frac{k^\gamma T}{n^\gamma}$, $n \geq 1$, $k = 0, 1, \dots, n$, and let $s_k = s_k^{(n)} = t_k - t_{k-1}$, $k = 1, \dots, n$. Then we have the following.

Theorem 3 Let $m \geq 1$ and $Q_{(s)}$, $s > 0$ be an m -similar family of linear operators in $C_b(\mathbf{R}^N)$. Then we have the following.

For $\gamma \in (0, m - 1)$, there is a constant $C > 0$ such that

$$\|P_T f - Q_{(s_n)} Q_{(s_{n-1})} \cdots Q_{(s_1)} f\|_\infty \leq C n^{-\gamma/2} \|\nabla f\|_\infty, \quad f \in C_b^\infty(\mathbf{R}^N), \quad n \geq 1.$$

For $\gamma = m - 1$, there is a constant $C > 0$ such that

$$\|P_T f - Q_{(s_n)} Q_{(s_{n-1})} \cdots Q_{(s_1)} f\|_\infty \leq C n^{-\frac{m-1}{2}} \log(n+1) \|\nabla f\|_\infty,$$

$$f \in C_b^\infty(\mathbf{R}^N), \quad n \geq 1.$$

For $\gamma > m - 1$, there is a constant $C > 0$ such that

$$\|P_T f - Q_{(s_n)} Q_{(s_{n-1})} \cdots Q_{(s_1)} f\|_\infty \leq C n^{-\frac{m-1}{2}}, \|\nabla f\|_\infty, \quad f \in C_b^\infty(\mathbf{R}^N), \quad n \geq 1.$$

Definition 4 We say that a $\mathcal{L}((A))$ -valued random variable Z is m - \mathcal{L} -moment similar, $m \geq 1$, if

$$E[\langle j_m(Z), j_m(Z) \rangle^n] < \infty \quad \text{for any } n \geq 1,$$

and if

$$E[j_m(\exp(Z))] = E[j_m(X(1))].$$

Theorem 5 Let $m \geq 1$ and Z be a $\mathcal{L}((A))$ -valued m - \mathcal{L} -moment similar random variable. Also, let $Y : (0, 1] \times \Omega \rightarrow C(\mathbf{R}^N; \mathbf{R}^N)$ be a measurable map such that

$$\sup_{s \in (0, 1], x \in \mathbf{R}^N} s^{-(m+1)/2} E[|Y(s)(x)|] < \infty$$

and

$$E[\sup_{|x| \leq n} |Y(s)(x)|] < \infty, \quad s \in (0, 1], \quad n \geq 1.$$

Let us define a linear map $Q_{(s)}$, $s > 0$, in $C_b(\mathbf{R}^N)$ by

$$(Q_{(s)} f)(x) = E[f(\exp(\Phi(j_m(\Psi_s(Z))))(x) + Y(s)(x))], \quad f \in C_b(\mathbf{R}^N).$$

Then $\{Q_{(s)}; s \in (0, 1]\}$ is m -similar.

We can show the following characterization theorem for 5- \mathcal{L} -moment similar random variables.

Theorem 6 Let Z be an $\mathcal{L}_{\leq 5}(A)$ -valued random variable. Then Z is 5- \mathcal{L} moment similar, if and only if there are random variables ξ_i , $i = 1, \dots, d$, η_{ij} , $1 \leq i < j \leq d$, $\zeta_{ij}^{(3)}$, $i, j = 1, \dots, d$, $i \neq j$, $\zeta_{0i}^{(4)}$, $i = 1, \dots, d$, and $\mathcal{L}_m(A)$ -valued random variables $\rho^{(m)}$, $m = 3, 4, 5$, satisfying the following.

$$(1) \quad Z = \sum_{i=1}^d \xi_i v_i + (v_0 + \sum_{1 \leq i < j \leq d} \eta_{ij}^{(2)} [v_i, v_j]) + (\sum_{1 \leq i \neq j \leq d} \zeta_{ij}^{(3)} [[v_i, v_j], v_j] + \rho^{(3)}) + (\sum_{i=1}^d \zeta_{0i}^{(4)} [[v_0, v_i], v_i] + \rho^{(4)}) + \rho^{(5)}.$$

$$(2) \quad E[\xi_i] = E[\xi_i^3] = E[\xi_i^5] = 0,$$

$$E[\xi_i^2] = 1, \quad E[\xi_i^4] = 3, \quad i = 1, \dots, d,$$

$$E[\eta_{ij}] = 0, \quad E[\eta_{ij}^2] = 1, \quad 1 \leq i < j \leq d,$$

$$E[\prod_{i=1}^d \xi_i^{\alpha_i} \prod_{1 \leq i < j \leq d} \eta_{ij}^{\beta_{ij}}] = \prod_{i=1}^d E[\xi_i^{\alpha_i}] \prod_{1 \leq i < j \leq d} E[\eta_{ij}^{\beta_{ij}}]$$

for any non-negative integers α_i , $i = 1, \dots, d$, and β_{ij} , $1 \leq i < j \leq d$ with $\sum_{i=1}^d \alpha_i + \sum_{1 \leq i < j \leq d} 2\beta_{ij} \leq 5$.

$$(3) \quad E[\zeta_{ij}^{(3)}] = 0, \quad E[\xi_k \zeta_{ij}^{(3)}] = \frac{1}{12} \delta_{ik} \text{ for any } 1 \leq i, j, k \leq d, i \neq j, \text{ and}$$

$$E[(\xi_i \xi_j \zeta_{k\ell}^{(3)})] = 0, \quad 1 \leq i, j, k, \ell \leq d, \quad k \neq \ell,$$

$$E[(\eta_{ij} \zeta_{k\ell}^{(3)})] = 0 \quad 1 \leq i, j, k, \ell \leq d, \quad i < j, \quad k \neq \ell,$$

and

$$E[(\prod_{i=1}^d \xi_i^{\alpha_i} \prod_{1 \leq i < j \leq d} \eta_{ij}^{\beta_{ij}}) \rho^{(3)}] = 0$$

for any non-negative integers α_i , $i = 1, \dots, d$, and β_{ij} , $1 \leq i < j \leq d$ with $\sum_{i=1}^d \alpha_i + \sum_{1 \leq i < j \leq d} 2\beta_{ij} \leq 2$.

$$(4) \quad E[\zeta_{0i}^{(4)}] = \frac{1}{12}, \quad E[\xi_j \zeta_{0i}^{(4)}] = 0, \quad 1 \leq i, j \leq d, \text{ and}$$

$$E[\rho^{(4)}] = E[\xi_i \rho^{(4)}] = 0, \quad i = 1, \dots, d.$$

$$(5) \quad E[\rho^{(5)}] = 0.$$

References

- [1] Kusuoka, S., Approximation of Expectation of diffusion processes and Mathematical Finance, Advanced Studies in Pure Mathematics 31, Proceedings of Final Taniguchi Symposium, Nara 1998,(edited by Sunada, T.), Mathematical Society of Japan, 2001, pp. 147-165.
- [2] Kusuoka, S., Malliavin Calculus revisited, to appear in J. Math. Sci. Univ. Tokyo.
- [3] Lyons, T., and N. Victoir, Cubature on Wiener Space, Preprint
- [4] Ninomiya, S., A New Simulation Scheme of Diffusion Processes: Application of the Kusuoka Approximation to Finance Problems, Mathematics and Computer in Simulation, to appear.
- [5] Reutenauer, C., Free Lie Algebras, Clarendon Press, Oxford, 1993.